Photomedicine and Laser Surgery Volume 32, Number 9, 2014 Mary Ann Liebert, Inc. Pp. 490–494

DOI: 10.1089/pho.2014.3715

## Effect of Diode Laser in the Treatment of Patients with Nonspecific Chronic Low Back Pain: A Randomized Controlled Trial

Francesco Vallone, MD,1 Stefano Benedicenti, DDS,2 Eugenio Sorrenti, DDS,2 Irene Schiavetti, MD,3 and Francesca Angiero, MD, DDS2 Abstract

Background data: Low back pain is a common, highly debilitating condition, whose severity is variable. This study evaluated the efficacy of treatment with Ga-Al-As diode laser (980 nm) with a large diameter spot (32 cm²), in association with exercise therapy, in reducing pain. Objective: The present study aimed to evaluate the pain reduction efficacy of treatment with the Ga-Al-As diode laser (980 nm) in combination with exercise therapy, in patients with chronic low back pain (CLBP).

Methods: This study evaluated 100 patients with CLBP (mean age 60 years) who were randomly assigned to two groups. The laser plus exercises group (Laser + EX: 50 patients) received low-level laser therapy (LLLT) with a diode laser, 980 nm, with a specific handpiece [32 cm² irradiation spot size, power 20Win continuous wave (CW), fluence 37.5J/cm², total energy per point 1200 J] thrice weekly, and followed a daily exercise schedule for 3 weeks (5 days/week). The exercises group (EX: 50 patients) received placebo laser therapy plus daily exercises. The outcome was evaluated on the visual analogue pain scale (VAS), before and after treatment.

Results: At the end of the 3 week period, the Laser +EX group showed a significantly greater decrease in pain than did the EX group. There was a significant difference between the two groups, with average D VAS scores of 3.96 (Laser +EX group) and 2.23 (EX group). The Student's t test demonstrated a statistically significant difference between the two groups, at p < 0.001.

Conclusions: This study demonstrated that the use of diode laser (980 nm) with large diameter spot size, in association with exercise therapy, appears to be effective. Such treatment might be considered a valid therapeutic option within rehabilitation programs for nonspecific CLBP.